

- ORZELL, S. L. 1983. Natural area inventory and floristic analysis of fens in selected southeastern Missouri counties. M. S. thesis, Southern Illinois University, Carbondale.
- _____. 1984. Natural area inventory and floristic analysis of fens in selected southeastern Missouri counties, Part II. *Missouriensis* 5:109 – 116.
- REZNICEK, A. A. and P. W. BALL. 1980. The taxonomy of *Carex* section *Stellulatae* in North America north of Mexico. *Contr. Univ. Mich. Herb.* 14:153 – 203.
- WHEELER, G. A. and G. B. OWNBEY. 1984. Annotated list of Minnesota Carices, with phytogeographical and ecological notes. *Rhodora* 86:151 – 231.

AGRIMONIA INCISA (ROSACEAE) NEW TO TEXAS.—In the course of routine identifications for Edward C. Fritz of the Texas Committee on Natural Resources, specimens of *Agrimonia incisa* T. & G. were identified. Steve Orzell (Texas Natural Heritage Program, Texas Parks and Wildlife Department; pers. comm.) did not find any specimens at TEX from Texas. From the literature, the species ranges from the Carolinas to northern Florida westward to Mississippi. While one would expect it to occur in Louisiana, no reports of this taxon appear in the literature from that state.

Jasper Co.: dry longleaf uplands; upper Trout Creek, 0.8 mi E of FR 330 on Plum Ridge Road and 0.5 mi S, 21 Aug 1988, *Edward C. Fritz & John Ward* 8884 (ASTC, SMU).

—*Wm. F. Mahler, SMU Herbarium/BRIT, Dallas, TX 75275, U.S.A.*

LEONURUS MARRUBIASTRUM L. (LAMIACEAE), NEW TO MISSOURI—*Leonurus marrubiastrum* L., Biennial Motherwort, a native to northern Asia and eastern Europe (Ball 1972), is one of the numerous European invaders to the eastern United States. It occurs in isolated sites from Delaware and eastern Pennsylvania to Illinois and south to Florida (Fernald 1950; Gleason 1952). In the central U.S., it is found near the Missouri River in only seven counties in four states: northeastern Kansas (Douglas County), in east central Nebraska (Washington and Douglas counties) and southeastern Nebraska (Richardson and Nemah counties), western Iowa (Harrison County), and southeastern South Dakota (Hutchinson County) (Gr. Plains Fl. Assoc. 1977).

Biennial Motherwort was collected for the first time in Missouri during the summers of 1986 and 1987 at the Jamerson McCormack Wildlife Refuge Area in Holt County. These collections were made at the edge of woods in the refuge's parking lot. It should now be expected to occur in